

Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP)

Summary

Code	001-403-005
Immunogen	Recombinant Fc region of mouse IgG2a
Host	Alpaca pacous
Isotype	VHH domain of alpaca IgG2b/2c fused to goat IgG Fc
Conjugate	HRP
Specificity	Mouse IgG2a (Fc γ fragment specific)
Cross-Reactivity	No cross-reactivity with mouse IgG1/2b/3, mouse IgM, rabbit, human, cynomolgus, rat, goat IgG
Purity	Recombinant Expression and Affinity purified
Concentration	1mg/ml
Formation	Liquid, 10mM PBS (pH 7.5), 0.05% sucrose, 0.1% trehalose, 0.01% proclin300, 50% Glycerol
Storage	Store at -20 °C(Avoid freeze / thaw cycles), protect from light, Stable for 12 months at -20°C

Description

Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP) is designed for detecting mouse IgG2a Fc γ fragment specifically. Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP) is based on monoclonal, recombinant, goat IgG Fc fused single domain antibody to mouse IgG2a Fc γ fragment coupled to HRP. Based on immunoelectrophoresis and/or ELISA, Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP) reacts with the Fc fragment of mouse IgG2a selectively, no reactivity with other mouse IgG subclasses, mouse IgM, or the Fab portion of mouse immunoglobulins.

Background

Most monoclonal antibodies are generated in mouse. There are five antibody isotypes (IgA, IgD, IgE, IgG, and IgM) from mouse. Each isotype has a different heavy chain. Mouse IgG constitutes 75% of serum immunoglobulins, and IgG is the predominant form of first antibody produced from mouse. Mouse IgG consists of five subclasses-IgG1, IgG2a, IgG2b, IgG2c(inbred mouse strains with the Igh1-b allele have IgG2c isotype instead of IgG2a), IgG3. They are highly homologous and differ mainly in the hinge region. The whole IgG molecule possesses both the Fc region and the Fab region, which possessing the epitope-recognition site. The IgG contains two heavy and light chains, and the heavy chain is about 50 KD and the light chain is about 25 KD. The common IgG is monomeric with a molecular weight of approximately 150 kD.

VHH are single-domain antibodies derived from the variable regions of heavy chain of Camelidae immunoglobulin. The size of VHH is extremely small(<15KDa) compared to other forms of antibody fragment, which significantly increase the permeability of VHH. Thus VHH is considered of great value for research, diagnostics and therapeutics.

Benefits


High lot-to-lot consistency
 Increased sensitivity and higher affinity
 Animal-free production

Suggested Working Concentration


ELISA	1:10000-1:50000
WB	1:10000-1:50000

Dilution factors are presented in the form of a range because the optimal dilution is a function of many factors, such as antigen density, permeability, etc. The actual dilution used must be determined empirically.

This product is for research use only and is not approved for use in humans or in clinical



A titer ELISA of mouse IgG2a. The plate was coated with different amounts of mouse IgG2a. 1:10000 dilution of Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP) was used as the detection antibody.



ELISA of specificity for different species of IgG. The plate was coated with 2ug/ml of different IgG. 1:1000 dilution of Anti-Mouse IgG2a(Fc γ Fragment specific), AlpHcAbs[®] Goat antibody(HRP) was used as the detection antibody.

This product is for research use only and is not approved for use in humans or in clinical